


Alain, C., radio R-II, et Karine Massicotte-Tisluck M.D. Faculté de Médecine et des sciences de la santé, Université de Sherbrooke

Introduction

Renseignements manquants ou incomplets; quelles sont les causes ?




Objectifs d'apprentissage


Évaluer l'impact des renseignements cliniques sur

- Exactitude de l'interprétation
- Niveau de confiance du radiologiste
- Pertinence du rapport
- Efficacité du radiologiste


Exactitude de l'interprétation




50 TDM variées interprétées par 2 radiologistes (total 100 rapports)⁵
 SANS puis AVEC renseignements cliniques




Niveau de confiance



Pertinence du rapport



Efficacité



Conclusion

Renseignements cliniques ↑ pertinence + exactitude + niveau de confiance du rapport
 Pistes de solutions pour obtenir renseignements complets et de qualité :

- Schéma/graphique indiquant site douleur
- Questionnaire rempli par patient ou technologue
- Formulaire/canevas spécifique lors prescription examen

Références

- Castillo, C., Steffens, T., Sim, L., & Caffery, L. (2021). The effect of clinical information on radiology reporting: A systematic review. *Journal of medical radiation sciences*, 68(1), 60-74.
- Lacson, R., Laroya, R., Wang, A., Kapoor, N., Glazer, D. I., Shinagare, A., Ip, I. K., Malhotra, S., Hentel, K., & Khorasani, R. (2018). Integrity of clinical information in computerized order requisitions for diagnostic imaging. *Journal of the American Medical Informatics Association : JAMIA*, 25(12), 1651-1656.
- Qureishi, A., Garas, G., Shah, J., & Birchall, J. (2014). A two-cycle prospective audit of temporal bone computed tomography scan requests: improving the clinical applicability of radiology reports. *The Journal of laryngology and otology*, 128(1), 49-52.
- Mullins, M. E., Lev, M. H., Schellingerhout, D., Koroshetz, W. J., & Gonzalez, R. G. (2002). Influence of availability of clinical history on detection of early stroke using unenhanced CT and diffusion-weighted MR imaging. *AJR. American journal of roentgenology*, 179(1), 223-228.
- Leslie, A., Jones, A. J., & Goddard, P. R. (2000). The influence of clinical information on the reporting of CT by radiologists. *The British journal of radiology*, 73(874), 1052-1055.
- Doshi, A. M., Huang, C., Ginocchio, L., Shanbhogue, K., & Rosenkrantz, A. B. (2017). Impact of patient questionnaires on completeness of clinical information and identification of causes of pain during outpatient abdominopelvic CT interpretation. *Abdominal radiology (New York)*, 42(12), 2946-2950.
- Cohen, M. D., & Ellett, M. (2012). Quality of communication: different patterns of reporting the location of the tip of a nasogastric tube. *Academic radiology*, 19(6), 651-653.
- Sarwar, A., Wu, J. S., Kung, J., Brook, A., Lee, K. S., Gauguet, J. M., & Rosen, M. P. (2014). Graphic representation of clinical symptoms: a tool for improving detection of subtle fractures on foot radiographs. *AJR. American journal of roentgenology*, 203(4), W429-W433.