

On Bayesian credible sets in restricted parameter space problems and lower bounds for frequentist coverage ¹

ÉRIC MARCHAND^a, WILLIAM E. STRAWDERMAN^b

a Université de Sherbrooke, Département de mathématiques, Sherbrooke Qc, CANADA, J1K 2R1

(e-mail: eric.marchand@usherbrooke.ca)

b Rutgers University, Department of Statistics and Biostatistics, 501 Hill Center, Busch Campus,

Piscataway, N.J., USA, 08855 (e-mail: straw@stat.rutgers.edu)

SUMMARY

For estimating a lower bounded parametric function in the framework of Marchand and Strawderman (2006), we provide through a unified approach a class of Bayesian confidence intervals with credibility $1 - \alpha$ and frequentist coverage probability bounded below by $\frac{1-\alpha}{1+\alpha}$. In cases where the underlying pivotal distribution is symmetric, the findings represent extensions with respect to the specification of the credible set achieved through the choice of a *spending function*, and include Marchand and Strawderman's HPD procedure result. For non-symmetric cases, the determination of a such a class of Bayesian credible sets fills a gap in the literature and includes an "equal-tails" modification of the HPD procedure. Several examples are presented demonstrating wide applicability.

AMS 2000 subject classifications: 62C10, 62F15, 62F25, 62F30

Key words and phrases: Bayesian methods, Credible sets, Frequentist coverage probability, Lower bound, Restricted Parameter, Spending function

¹December 17, 2012

1. Introduction

Bayesian credible sets are not designed (e.g., Robert, 2011) and are far from guaranteed (Fraser, 2011) to have satisfactory, exact or precise frequentist coverage but it is nevertheless of interest to investigate (Wasserman, 2011) to what extent there is convergence or divergence in various situations. A historically resonating example where there is exact convergence arises for estimating the mean of a $N(\mu, \sigma^2)$ distribution, and where the use of the non-informative prior leads to a $(1 - \alpha) \times 100\%$ HPD credible set (i.e. the z or t confidence interval) with exact frequentist coverage. This, however, is very much the exception. Even, in the simple presence of a lower bound on the mean parameter μ (e.g., Mandelkern, 2002), with the prior taken to be the truncation of the non-informative prior onto the restricted parameter space, the frequentist coverage of the $(1 - \alpha) \times 100\%$ HPD credible set fluctuates from its credibility (or nominal coverage) $1 - \alpha$. However, the HPD procedure does not fare poorly as a frequentist procedure for large $1 - \alpha$ as witnessed by the lower bound $\frac{1-\alpha}{1+\alpha}$ on its frequentist coverage due to Roe and Woodroffe (2000, known σ^2) and Zhang and Woodroffe (2003, unknown σ^2), as well as the better lower bound $1 - \frac{3\alpha}{2}$ (for $\alpha < 1/3$, known σ^2) obtained by Marchand et al. (2008).

In a generalization of the above, Marchand and Strawderman (MS 2006) introduced a unified framework for which the $(1 - \alpha) \times 100\%$ HPD credible set of a lower bounded parametric function has frequentist coverage greater than $\frac{1-\alpha}{1+\alpha}$ for all values lying in the restricted parameter space. This framework, as well as its various applications, will be revisited in Sections 2 and 5, but let us consider for sake of illustration the basic examples: (i) $X \sim f_0(x - \theta)$ with known f_0 , $\theta \geq 0$; and (ii) $X \sim \text{Gamma}(\alpha, \theta)$ with $\theta \geq 1$ with known α . For location family densities as in (i) with f_0 unimodal and symmetric, Marchand and Strawderman's results apply for the flat prior on the truncated parameter space $[0, \infty)$ and the corresponding $(1 - \alpha) \times 100\%$ HPD credible set,

with the guarantee that the actual frequentist coverage is bounded below by $\frac{1-\alpha}{1+\alpha}$ for all $\theta \geq 0$. However, if f_0 is not symmetric, such a result does not hold in general (MS 2006, Example 1). The same is true for a vast number of so-called non-symmetric situations arising in Marchand and Strawderman’s framework, including the Gamma models in (ii) where the prior is given by $\frac{1}{\theta}\mathbb{I}_{[1,\infty)}(\theta)$, that is the truncation on $[1, \infty)$ of the usual non-informative prior $\frac{1}{\theta}\mathbb{I}_{(0,\infty)}(\theta)$. It is true that the bound holds for certain specific classes of f_0 ’s (MS 2006, Theorem 2, a), and it is also the case that numerical evaluations of a theoretical and unexplicit lower bound for frequentist coverage provides further evidence for satisfactory coverage for a specific Gamma model in (ii) (MS 2006, Example 2). Nevertheless, a clear analytical result or lower bound for frequentist coverage in such non-symmetric cases is lacking, and it our motivation here to try to fill this gap.

For a large variety of situations with a lower bounded parametric constraint, we obtain here a class of Bayesian $(1 - \alpha) \times 100\%$ credible sets which provide minimal frequentist probability coverage exceeding $\frac{1-\alpha}{1+\alpha}$. These Bayesian confidence intervals include an “equal-tails” modification, or approximation, of the HPD credible set, which also coincides with the latter in situations of underlying symmetry. Our findings are achieved by introducing and exploiting a *spending* function interpretation of Bayesian confidence intervals, and lead to a class of procedures (rather than a single one) which share the above lower bound for frequentist coverage. The rest of the paper is organized as follows. Preliminary results, definitions and model assumptions, including those related to the spending function associated with a Bayesian credible interval, are presented in Section 2, while Bayesian credible interval representations are outlined in Section 3. The main findings concerning frequentist coverage appear in Section 4 and various applications are presented and commented on in Section 5.

2. Definitions and preliminary results

2.1. Assumptions, invariance, pivot, prior, and implications

As in basic examples (i) and (ii), we consider model densities $f(x; \theta)$; $x \in \mathcal{X}$, $\theta \in \Theta \subset \mathbb{R}^p$; for an observable X , and we are concerned with interval estimation of a parametric function $\tau(\theta)$ ($\mathbb{R}^p \rightarrow \mathbb{R}$) with the additional constraint $\tau(\theta) \geq 0$. We assume there exists a pivot of the form $T(X, \theta) = \frac{a_1(X) - \tau(\theta)}{a_2(X)}$; $a_2(\cdot) > 0$; such that $-T(X, \theta)$ has cdf G and Lebesgue density g_0 . This pivot assumption means that the frequentist or conditional distribution of $T(X, \theta)$, or $-T(X, \theta)$, given θ does not depend on θ , $\theta \in \mathbb{R}^p$. We can thus set G as the common cdf of $-T(X, \theta)$. In the basic location-family example (i) with $X \sim f_0(x - \theta)$ ($= g_0(\theta - x)$, say), the above is illustrated by the fact that $-T(X, \theta) = \theta - X$ is a pivot with cdf G and pdf g_0 . In the Gamma example, or more generally scale families with $X \sim \frac{1}{\theta} f_1(\frac{x}{\theta})$, $\theta \geq 1$, a corresponding $T(X, \theta)$ pivot is obtained with $a_1(X) = \log(X)$, $a_2(X) = 1$, $\tau(\theta) = \log(\theta)$.

We further assume that the unrestricted decision problem is invariant under a group \mathcal{G} of transformations and that the pivot satisfies the invariance requirement $T(x, \theta) = T(gx, \bar{g}\theta)$, for all $x \in \mathcal{X}$, $\theta \in \Theta$, $g \in \mathcal{G}$, $\bar{g} \in \bar{\mathcal{G}}$, with \mathcal{X} , Θ , G , and \bar{G} being isomorphic. For instance, in basic example (i), the invariance is achieved with the additive group G on \mathbb{R}^p and since $T(x, \theta) = x - \theta = (x + g) - (\theta + g) = T(gx, \bar{g}\theta)$ for all group elements g .

Collecting the above assumptions, we have for further reference.

Assumption 1. *We have a model density $f(x; \theta)$; $x \in \mathcal{X}$, $\theta \in \Theta$; for an observable X , with both X and θ being vectors, and we seek to estimate a parametric function $\tau(\theta)$ ($\mathbb{R}^p \rightarrow \mathbb{R}$) with the constraint $\tau(\theta) \geq 0$. We assume there exists a pivot $T(X, \theta) = \frac{a_1(X) - \tau(\theta)}{a_2(X)}$; $a_2(\cdot) > 0$; such that $-T(X, \theta)$ has cdf G and Lebesgue density g_0 . We further assume that the decision problem is*

invariant under a group \mathcal{G} of transformations and that the pivot satisfies the invariance requirement $T(x, \theta) = T(gx, \bar{g}\theta)$, for all $x \in \mathcal{X}$, $\theta \in \Theta$, $g \in \mathcal{G}$, $\bar{g} \in \bar{\mathcal{G}}$, with \mathcal{X} , Θ , G , and \bar{G} being isomorphic.

We consider prior measures π_H and π_0 , where $\pi_0(\theta) = \pi_H(\theta)I_{[0,\infty)}(\tau(\theta))$, and π_H is the Haar right invariant measure which satisfies the property $\pi_H(A\bar{g}) = \pi_H(A)$ for every measurable subset A of Θ , and for every $g \in G$. The right Haar measure π_H exists and is unique up to a multiplicative constant for locally compact groups such as location, scale, and location-scale. For the basic location and the Gamma model (or scale model) examples of the Introduction, right Haar invariant measures are given by $\pi_H(\theta) = 1$ and $\pi_H(\theta) = \frac{1}{\theta}$ respectively. For a sample from a location-scale family with $X_i \sim \text{ind. } \frac{1}{\theta_2} f_2(\frac{x_i - \theta_1}{\theta_2})$, $i = 1, \dots, n$, the common non-informative prior $\pi(\theta) = \frac{1}{\theta_2}$ is right Haar invariant. We refer to Berger (1985) or Eaton (1989) for detailed treatments of invariance and Haar invariant measures.

A key feature relative to Assumption 1 and the choice of the right Haar invariant measure is that the frequentist distribution of $T(X, \theta)$; which is free of θ by virtue of the pivot assumption for $T(X, \theta)$; coincides with the posterior distribution of $T(x, \theta)$ under π_H for any given x , i.e.,

$$T(x, \theta)|x =^d T(X, \theta)|\theta, \text{ for all } x, \theta. \quad (1)$$

We will pursue, after the next Lemma, by illustrating the above and drawing implications of immediate interest. For sake of completeness, we reproduce here a key lemma from MS(2006) justifying (1) and we refer to their work for further details.

Lemma 1. (MS, 2006, Corollary 1) Suppose \mathcal{X} , Θ , G , and \bar{G} are all isomorphic, and that $T(X, \theta)$ is a function for which $T(x, \theta) = T(gx, \bar{g}\theta)$, for all $x \in \mathcal{X}$, $\theta \in \Theta$, $g \in G$, $\bar{g} \in \bar{G}$. Then condition (1) holds, that is $P_\theta[T(X, \theta) \in B] = P^{\pi_H(\theta|x)}[T(X, \theta) \in B]$ for each measurable set B .

Now, for the basic unrestricted location family example with the flat prior $\pi_H(\theta) = 1$, which is Haar

right invariant, observe that the posterior density of θ is given by

$$\pi_H(\theta|x) = \frac{f_0(x - \theta)}{\int_{\theta} f_0(x - \theta) d\theta} = f_0(x - \theta) = g_0(\theta - x),$$

so that the posterior density of $-T(x, \theta) = \theta - x$ associated with π_H is given by g_0 as well. This correspondence for basic example (i) illustrates property (1) which is, of course, more general under Assumption 1.

In general, observe that the posterior cdf under π_H for $\tau(\theta)$ is available from the fact that $-T(X, \theta) = \frac{\tau(\theta) - a_1(X)}{a_2(X)} \sim G$ yielding

$$P_{\pi_H}(\tau(\theta) \leq y|x) = G\left(\frac{y - a_1(x)}{a_2(x)}\right). \quad (2)$$

Now, under the truncation π_0 of π_H , the above correspondence between the frequentist and posterior distributions of $-T(X, \theta)$ does not hold, and the posterior cdf under π_0 of $\tau(\theta)$ differs. However, we can still express the posterior distribution of $\tau(\theta)$ under π_0 in terms of π_H and G . Indeed, with $\pi_0(\theta) = \pi_H(\theta) \mathbb{I}_{[0, \infty)}(\tau(\theta))$ and $\frac{\tau(\theta) - a_1(x)}{a_2(x)}|x \sim G$ under π_H , we have for a measurable set $A \subset \Theta_0 = \{\theta \in \Theta : \tau(\theta) \geq 0\}$, and for any x :

$$\begin{aligned} P_{\pi_0}(\theta \in A|x) &= \int_A \pi_0(\theta|x) d\theta \\ &= \frac{\int_A \pi_0(\theta) f(x|\theta) d\theta}{\int_{\Theta_0} \pi_0(\theta) f(x|\theta) d\theta} \\ &= \frac{\int_A \pi_H(\theta) f(x|\theta) d\theta}{\int_{\Theta_0} \pi_H(\theta) f(x|\theta) d\theta} \\ &= \frac{P_{\pi_H}(\theta \in A|x)}{P_{\pi_H}(\theta \in \Theta_0|x)}. \end{aligned}$$

In terms of the posterior survival function of $\tau(\theta)$ under π_0 , the above yields along with (2), for $y \geq 0$,

$$P_{\pi_0}(\tau(\theta) \geq y|x) = \frac{P_{\pi}(\tau(\theta) \geq y|x)}{P_{\pi}(\tau(\theta) \geq 0|x)} = \frac{1 - G\left(\frac{y - a_1(x)}{a_2(x)}\right)}{1 - G\left(\frac{-a_1(x)}{a_2(x)}\right)}. \quad (3)$$

We will make use, in Section 3, of the above in setting and describing the bounds of Bayesian credible sets for $\tau(\theta)$ under π_0 .

2.2. The spending function associated with a Bayesian credible set

With the objective of constructing a $(1-\alpha) \times 100\%$ Bayesian credible set or region, the determination of a posterior distribution for $\tau(\theta)$ supported on $[0, \infty)$ leaves open many choices and various different approaches (e.g., Berger, 1985, section 4.3.2). The HPD credible set is one such region chosen to minimize volume and leading to intervals for unimodal posterior densities. In our set-up, $(1-\alpha) \times 100\%$ Bayesian credible intervals are, more generally, of the form $[l(x), u(x)]$, $x \in \mathbb{R}$, where $P(l(x) \leq \tau(\theta) \leq u(x)|x) = 1-\alpha$. An alternative (and equivalent) way to set or view the bounds $l(x)$ and $u(x)$, for a given x , is to focus on the complementary set $[0, l(x)) \cup (u(x), \infty)$ and to allocate (or “spend”) probabilities $\alpha - \alpha(x)$ and $\alpha(x)$ respectively on its two disjoint parts, with $\alpha(x) \in [0, \alpha]$. It is clear (when the posterior density is absolutely continuous) that the choice $\alpha(x)$ leads to a unique choice of $[l(x), u(x)]$, and vice-versa. Since we are interested in the frequentist properties of such Bayesian credible intervals, we will represent this allocation as a *spending* function. Moreover, our findings guaranteeing minimal frequentist coverage of at least $\frac{1-\alpha}{1+\alpha}$ for a class of Bayesian credible sets will be conveniently expressed as conditions on the corresponding spending function.

Definition 1. *For a given prior π for θ and a credibility coefficient $1 - \alpha$, a spending function $\alpha(\cdot) : \mathbb{R}^p \rightarrow [0, \alpha]$ is a function such that, for all x , $P_\pi(\tau(\theta) \geq u(x)|x) = \alpha(x)$, $P_\pi(\tau(\theta) \leq l(x)|x) = \alpha - \alpha(x)$, and $[l(x), u(x)]$ is a $(1 - \alpha) \times 100\%$ Bayesian credible interval for $\tau(\theta)$.*

For example, a lower-tailed credible interval for a given x corresponds to the selection $\alpha(x) = \alpha$, an upper tailed credible interval corresponds to $\alpha(x) = 0$, and an equal tailed (based on the posterior π) corresponds to $\alpha(x) = \alpha/2$.

2.3. Checklist

To facilitate the further presentation of the results, here is a list of definitions and notations used.

Checklist

- $1 - \alpha$: credibility or posterior coverage or nominal frequentist coverage ($\alpha \in (0, 1)$)
- $T(X, \theta) = \frac{a_1(X) - \tau(\theta)}{a_2(X)}$: pivot
- π_H : unrestricted prior density chosen as the right Haar invariant measure
- π_0 : prior density given by the truncation of π_H onto the restricted parameter space
- G : cumulative distribution function (cdf) of $-T(X, \theta)|x$ and of $-T(X, \theta)|\theta$ under π_H (which coincide for all x, θ)
- $g_0 = G'$: probability density function (pdf) of $-T(X, \theta)$
- G^{-1} : inverse cdf
- $\alpha(\cdot)$: spending function
- $I_{\pi_0, \alpha(\cdot)}(X) = [l(X), u(X)]$: Bayesian credible set of credibility $1 - \alpha$ associated with the prior π_0 and the spending function $\alpha(\cdot)$
- $C(\theta)$: the frequentist coverage at θ of the confidence interval $I_{\pi_0, \alpha(\cdot)}(X)$ given by $C(\theta) = P_\theta(I_{\pi_0, \alpha(\cdot)}(X) \ni \tau(\theta))$
- $y_0 = -G^{-1}(\frac{\alpha}{1+\alpha})$
- $t(x) = \frac{a_1(x)}{a_2(x)}$
- $\Delta_0(x) = (1 - \alpha)(1 - G(-t(x)))$

3. Bayesian credible intervals: representations and properties

In this section, we expand upon two different, yet equivalent, and instructive approaches to constructing a credible set for $\tau(\theta)$ associated with prior π_0 . These are: (A) the spending function approach, and (B) the approach based on the quantiles of the pivot.

(A) (Spending function approach)

As seen above, a $(1 - \alpha) \times 100\%$ credible interval for $\tau(\theta)$ associated with prior π_0 can be generated by a *spending* function $\alpha(\cdot) : \mathbb{R}^p \rightarrow [0, \alpha]$, such that $I_{\pi_0, \alpha(\cdot)}(X) = [l(X), u(X)]$ with $P_{\pi_0}(\tau(\theta) \geq u(x)|x) = \alpha(x)$. More precisely, we have the following under Assumption 1.

Lemma 2. *For a given spending function $\alpha(\cdot)$, the bounds of $I_{\pi_0, \alpha(\cdot)}(x)$ are given by: $l_{\alpha(\cdot)}(x) = a_1(x) + a_2(x)G^{-1}\{G(-t(x)) + (\alpha - \alpha(x))(1 - G(-t(x)))\}$ and $u_{\alpha(\cdot)}(x) = a_1(x) + a_2(x)G^{-1}\{1 - \alpha(x)(1 - G(-t(x)))\}$, with $t(x) = \frac{a_1(x)}{a_2(x)}$.*

Proof. With the survival function $P_{\pi_0}(\tau(\theta) \geq y|x) = \frac{1 - G(\frac{y - a_1(x)}{a_2(x)})}{1 - G(-t(x))}$, as given in (3), we obtain for $\beta \in (0, 1), y > 0$, $P_{\pi_0}(\tau(\theta) \geq y|x) = \beta \Leftrightarrow y = a_1(x) + a_2(x)G^{-1}\{1 - \beta + \beta G(-t(x))\}$, and the result follows with the choices $\beta = \alpha(x)$ and $\beta = 1 - (\alpha - \alpha(x))$ for $u(x)$ and $l(x)$ respectively. \square

Example 1. *The HPD procedures studied by MS (2006) for symmetric about 0 and unimodal g_0 are given by the bounds $l(x) = \max\{0, a_1(x) + a_2(x)G^{-1}(\frac{1 - (1 - \alpha)G(t(x))}{2})\}$ and $u(x) = a_1(x) + a_2(x) \min\{G^{-1}(1 - \alpha G(t(x))), G^{-1}(\frac{1 + (1 - \alpha)G(t(x))}{2})\}$. With these given bounds, one may verify directly from (3) that the corresponding spending function is equal to*

$$\min\left\{\alpha, \frac{\alpha}{2} + \frac{G(-t(x))}{2(1 - G(-t(x)))}\right\}, \quad (4)$$

with $\alpha(x) = \alpha$ if and only if $t(x) \leq -G^{-1}(\frac{\alpha}{1 + \alpha}) = G^{-1}(\frac{1}{1 + \alpha})$ since g_0 is symmetric about 0. Conversely, applying Lemma 2 with the spending function choice $\alpha(\cdot)$ in (4) leads to the HPD procedure above (using the equality of $G(\cdot)$ and $1 - G(-\cdot)$ for symmetric about 0 g_0 's).

(B) (Approach based on quantiles of the pivot)

Alternatively, a second approach for cases where $l(x) > 0$ begins with choices γ_1 and γ_2 , which

will be made for each x , such that $G(\gamma_2) - G(-\gamma_1) = \Delta$, for a given $\Delta \in (0, 1)$. Since, for any x , we require $1 - \alpha = P_{\pi_0}(l(x) \leq \tau(\theta) \leq u(x)|x)$, we must have by (3):

$$G\left(\frac{u(x) - a_1(x)}{a_2(x)}\right) - G\left(\frac{l(x) - a_1(x)}{a_2(x)}\right) = (1 - \alpha)(1 - G(-t(x))),$$

and this can be achieved with choices $-\gamma_1$ and γ_2 above for $\Delta = \Delta_0(x) = (1 - \alpha)(1 - G(-t(x)))$ yielding $\frac{u(x) - a_1(x)}{a_2(x)} = \gamma_2(\Delta_0(x))$ and $\frac{l(x) - a_1(x)}{a_2(x)} = -\gamma_1(\Delta_0(x))$, in other words

$$l(x) = a_1(x) - a_2(x) \gamma_1(\Delta_0(x)), \text{ and } u(x) = a_1(x) + a_2(x) \gamma_2(\Delta_0(x)), \quad (5)$$

whenever $l(x) > 0$. In view of the lower bound restriction on $\tau(\theta)$ (i.e., $\tau(\theta) \geq 0$), and the corresponding requirement that $l(X) \geq 0$, observe that not all choices of $-\gamma_1$ (and hence of γ_2) are feasible in (5) and that we must have

$$-\gamma_1(\Delta_0(x)) \geq -\frac{a_1(x)}{a_2(x)}.$$

Example 2. *With the above construction in (5), an equal-tails choice of $-\gamma_1$ and γ_2 , that is $-\gamma_1(\Delta) = G^{-1}(\frac{1-\Delta}{2})$ and $\gamma_2(\Delta) = G^{-1}(\frac{1+\Delta}{2})$, leads to the credible interval bounds*

$$l(x) = a_1(x) - a_2(x) G^{-1}\left(\frac{1 - \Delta_0(x)}{2}\right), \text{ and } u(x) = a_1(x) + a_2(x) G^{-1}\left(\frac{1 + \Delta_0(x)}{2}\right), \quad (6)$$

when $l(x) > 0$. These above bounds coincide with those of the HPD procedure (when $l(x) > 0$) in the symmetric case of Example 1, as well as the spending function given in (4) as can be verified directly from (3).

NOTE: We wish to emphasize that the terminology “equal tails” does not mean $\alpha(x) = \alpha/2$ (i.e., equal tails under the posterior distribution), but rather refers to the choice of (equal tails) quantiles $-\gamma_1$ and γ_2 under G .

The next section’s lower bound of $\frac{1-\alpha}{1+\alpha}$ on frequentist coverage applies to a class of Bayesian credible intervals. This class will include an *equal-tails* credible interval $I_{\pi_0, \alpha_{eqt}(\cdot)}$ which relates to both

approaches presented in this section. On one hand, it borrows the bounds (and hence the spending function) of the HPD procedure for symmetric about 0 unimodal densities and, on the other hand, it is defined through the above equal-tailed choice (whenever $l(x) > 0$).

Definition 2. *In the context of Assumption 1, the G -equal-tails credible interval $I_{\pi_0, \alpha_{eqt}(\cdot)}(X)$ is given by the bounds $l(x) = \max\{0, a_1(x) + a_2(x)G^{-1}(\frac{1-(1-\alpha)G(t(x))}{2})\}$ and $u(x) = a_1(x) + a_2(x) \min\{G^{-1}(1-\alpha G(t(x))), G^{-1}(\frac{1+(1-\alpha)G(t(x))}{2})\}$. Equivalently, $I_{\pi_0, \alpha_{eqt}(\cdot)}(X)$ is given by the spending function*

$$\alpha_{eqt}(x) = \min\left\{\alpha, \frac{\alpha}{2} + \frac{G(-t(x))}{2(1-G(-t(x)))}\right\}. \quad (7)$$

4. Frequentist coverage properties

We study here the frequentist coverage properties, under Assumption 1, of a class of Bayesian credible intervals which includes the equal-tails credible interval $I_{\pi_0, \alpha_{eqt}(\cdot)}(X)$. This procedure, as well as Example 1's HPD procedure for symmetric g_0 , produces estimates of the form $[0, u(x)]$ if and only if $t(x) \leq y_0$, where $y_0 = -G^{-1}(\frac{\alpha}{1+\alpha})$ (and $t(x) = \frac{a_1(x)}{a_2(x)}$ as above). We thus focus on a class of credible intervals with the same behaviour. Said otherwise in terms of the spending function, we impose the choice $\alpha(x) = \alpha$ whenever $t(x) \leq y_0$. We hence seek conditions on $\alpha(x)$, for those x 's such that $t(x) \geq y_0$, for which minimal frequentist coverage is bounded below by $\frac{1-\alpha}{1+\alpha}$.

Theorem 1. *Under the conditions of Theorem 1 of Marchand and Strawderman (2006), that is Assumption 1, consider Bayesian credible intervals $I_{\pi_0, \alpha(\cdot)}$ associated with prior π_0 and a spending function $\alpha(\cdot)$ such that $\alpha(x) = \alpha$ for all x with $t(x) \leq y_0$. For the frequentist coverage $C(\theta) = P_\theta(I_{\pi_0, \alpha(\cdot)}(X) \ni \tau(\theta))$, we then have*

$$(a) \ C(\theta) = \frac{1}{1+\alpha} (> \frac{1-\alpha}{1+\alpha}) \text{ for all } \theta \text{ such that } \tau(\theta) = 0;$$

(b) Moreover, we have $C(\theta) > \frac{1-\alpha}{1+\alpha}$ for all θ such that $\tau(\theta) \geq 0$ as long as $\alpha(x)$ satisfies, for all x ,

$$\frac{(1-\alpha)G(-t(x)) + \frac{\alpha^2}{1+\alpha}}{1-G(-t(x))} \leq \alpha(x) \leq \left(\frac{\alpha}{1+\alpha}\right) \frac{1}{1-G(-t(x))}. \quad (8)$$

Proof.

(a) First, observe that for θ such that $\tau(\theta) = 0$, the pivot assumption for $-T(X, \theta) = \frac{\tau(\theta) - a_1(X)}{a_2(X)}$ implies that $-t(X) = -\frac{a_1(X)}{a_2(X)}$ has cdf G whenever $\tau(\theta) = 0$. Hence, for θ such that $\tau(\theta) = 0$, we have

$$P_\theta(I_{\pi_0, \alpha(\cdot)}(X) \ni 0) = P_\theta(\alpha(X) = \alpha) = P_\theta(t(X) \leq y_0) = 1 - G(-y_0) = \frac{1}{1+\alpha}.$$

(b) With the case $\tau(\theta) = 0$ addressed in part (a), we consider $\tau(\theta) > 0$. First, observe that the confidence interval $I_1(X) = [l_1(X), u_1(X)] = \max\{0, a_1(X) + a_2(X)G^{-1}(\frac{\alpha}{1+\alpha})\}, a_1(X) + a_2(X)G^{-1}(\frac{1}{1+\alpha})\}$ has the same frequentist coverage as $I_1^*(X) = [a_1(X) + a_2(X)G^{-1}(\frac{\alpha}{1+\alpha})\}, a_1(X) + a_2(X)G^{-1}(\frac{1}{1+\alpha})\}$ equal to $P_\theta(G^{-1}(\frac{\alpha}{1+\alpha}) \leq \frac{\tau(\theta) - a_1(X)}{a_2(X)} \leq G^{-1}(\frac{1}{1+\alpha})) = G(G^{-1}(\frac{1}{1+\alpha})) - G(G^{-1}(\frac{\alpha}{1+\alpha})) = \frac{1-\alpha}{1+\alpha}$. Now, we show that the given conditions on $\alpha(\cdot)$ imply that $I_{\pi_0, \alpha(\cdot)} \supseteq I_1$; with the inclusion being strict with probability greater than 0 for all θ ; which will lead to the result directly. Indeed, we have by the upper bound in (8) and Lemma 2: $u_{\alpha(\cdot)}(x) \geq a_1(x) + a_2(x)G^{-1}(1 - \frac{\alpha}{1+\alpha}) = u_1(x)$. Similarly, from the lower bound (8) and Lemma 2 we obtain $l(x) \leq a_1(x) + a_2(x)G^{-1}\{G(-t(x)) + \alpha(1 - G(-t(x))) - \frac{\alpha^2}{1+\alpha} - (1 - \alpha)G(-t(x))\} = a_1(x) + a_2(x)G^{-1}(\frac{\alpha}{1+\alpha}) = l_1(x)$. \square

Corollary 1. Under Assumption 1, the G -equal-tails credible interval $I_{\pi_0, \alpha(\cdot)}$, given in Definition 7, has minimum frequentist coverage $C(\theta)$ greater than $\frac{1-\alpha}{1+\alpha}$ for all θ such that $\tau(\theta) \geq 0$.

Proof. It suffices to show directly that (8) is satisfied for the selection $\alpha(x) = \alpha_{\text{eqt}}(x)$ given in (4) for x such that $t(x) \geq y_0$. Indeed, we have for such x 's:

$$\alpha_{\text{eqt}}(x)(1 - G(-t(x))) = \frac{\alpha}{2} + \frac{1 - \alpha}{2} G(-t(x)) \leq \frac{\alpha}{2} + \frac{1 - \alpha}{2} G(-y_0) = \frac{\alpha}{1 + \alpha},$$

and

$$\begin{aligned} \alpha_{\text{eqt}}(x)(1 - G(-t(x))) - (1 - \alpha) G(-t(x)) - \frac{\alpha^2}{1 + \alpha} &= \frac{\alpha(1 - \alpha)}{2(1 + \alpha)} - \frac{1 - \alpha}{2} G(-t(x)) \\ &\geq \frac{\alpha(1 - \alpha)}{2(1 + \alpha)} - \frac{1 - \alpha}{2} G(-y_0) = 0. \quad \square \end{aligned}$$

Remark 1. *In cases where the underlying pivotal distribution is non-symmetric, Corollary 1 is a new result, generalizing Theorem 1 of MS (2006), and is widely applicable given the lack of assumptions on g_0 . Also, the bounds of the equal-tails procedure are easier to evaluate than that of the HPD credible interval. And the findings of Theorem 1 go beyond a single procedure, even in the symmetric case, by providing a class of credible sets, as specified by a spending function, with frequentist coverage bounded below by $\frac{1-\alpha}{1+\alpha}$.*

We do not have a recommended prescription for the choice of the spending function among those specified by Theorem 1 as guaranteeing minimal frequentist coverage of at least $\frac{1-\alpha}{1+\alpha}$. The G -equal-tails choice is simple, intuitively appealing and matches the HPD procedure under symmetry of the pivotal density, while upper tailed and lower tailed choices are not allowed for x such that $t(x) \geq y_0$. The bounds in (8): (i) $\alpha_1(x) = \frac{(1-\alpha)G(-t(x)) + \frac{\alpha^2}{1+\alpha}}{1-G(-t(x))}$ and (ii) $\alpha_2(x) = \left(\frac{\alpha}{1+\alpha}\right) \frac{1}{1-G(-t(x))}$, are other interesting choices which push extremally $I_{\pi, \alpha(\cdot)}$ towards $+\infty$ and 0 respectively. Finally, along with these choices, it might be feasible to minimize the length of the credible interval under the restrictions imposed by Theorem 1.

5. Examples

At the risk of some redundancy with the examples provided by MS (2006), it is still beneficial here to present various applications with accompanying commentary. Assumption 1 is satisfied in all of the examples below with the underlying family of transformations (distributions) being either the location family, the scale family, or the location-scale family. In all of the examples, Theorem 1 and Corollary 1 provide conditions on the spending function $\alpha(\cdot)$ so that the Bayesian intervals $I_{\pi_0, \alpha(\cdot)}(X)$ have minimal frequentist coverage greater than $\frac{1-\alpha}{1+\alpha}$ for all θ such that $\tau(\theta) \geq 0$. These intervals include the equal-tails procedure given in Definition 2 and can be evaluated in general using the expression given in Lemma 2.

(A) (**a single location parameter**) $X \sim f_0(x - \theta)$; $\tau(\theta) = \theta \geq 0$; $T(X, \theta) = X - \theta$; $\pi_H(\theta) = \mathbb{I}_{\mathbb{R}}(\theta)$, $\pi_0(\theta) = \mathbb{I}_{[0, \infty)}(\theta)$. In such cases, all Bayes credible sets $I_{\pi_0, \alpha(\cdot)}$ (with credibility $1 - \alpha$), with the spending function $\alpha(\cdot)$ satisfying the conditions of Theorem 1 and the bounds in (8), have necessarily minimum frequentist coverage bounded below by $\frac{1-\alpha}{1+\alpha}$. Through the transformations $X \rightarrow X - a$ and $X \rightarrow -X + a$, one can reduce all lower bounded restrictions $\theta \geq a$ and upper bounded restrictions $\theta \leq a$ to the case $\theta \geq 0$ considered here and we will not make further explicit mention of such transformations below.

Remark 2. *Results such as those in (A) are applicable as well for several observations by conditioning on a maximal invariant statistic V . Such a maximal invariant statistic V is an ancillary statistic and specifically an invariant function such that every other invariant statistic is a function of V . Indeed, suppose that $X = (X_1, \dots, X_n) \sim f_0(x_1 - \theta, \dots, x_n - \theta)$, where f_0 is known and where the X_i 's are not necessarily independently distributed. Here, $V = (X_2 - X_1, \dots, X_n - X_1)$ is a maximal invariant statistic. One can then proceed, for a given value v of V , with an interval estimate*

$I_{\pi_0, \alpha(\cdot, v)}(X_1, v)$ as given in Lemma 2 with $G \equiv G_v$ representing the cdf of the pivot $X_1 - \theta$ conditional on $V = v$, and $\alpha(x, v)$ satisfying the conditions of Theorem 1 and (8). This is feasible by the pivot and ancillarity property with the joint distribution of $(X_1 - \theta, V)$ independent of θ . In such a case, Theorem 1 applies to the conditional frequentist coverage $C(\theta, v) = P_\theta(I_{\pi_0, \alpha(\cdot, v)}(X, v) \ni \tau(\theta) | V = v)$ yielding the inequality $C(\theta, v) > \frac{1-\alpha}{1+\alpha}$ for all $\theta \geq 0$. Since this is true for all v , the unconditional frequentist coverage $C(\theta)$ of the Bayes credible set $I_{\pi_0, \alpha(\cdot, \cdot)}(X, V)$ will also exceed $\frac{1-\alpha}{1+\alpha}$ for all $\theta \geq 0$ (see MS 2006, for more details related to a multivariate Student model). In the same vein, all the scenarios below (B to G), although presented for simplicity in the single observation case, are also applicable in presence of a sample by conditioning on a maximal invariant statistic.

(B) **(a lower bounded scale parameter)** $X \sim \frac{1}{\theta} f_1(\frac{x}{\theta}) \mathbb{I}_{(0, \infty)}(x)$ with $\theta \geq a$; $\tau(\theta) = \log(\theta) - \log(a) \geq 0$; $T(X, \theta) = \log(X) - \log(a) - \tau(\theta)$; $\pi_H(\theta) = \frac{1}{\theta} \mathbb{I}_{(0, \infty)}(\theta)$, $\pi_0(\theta) = \frac{1}{\theta} \mathbb{I}_{[0, \infty)}(\tau(\theta))$. Here, an interval estimate of $\tau(\theta)$ provides an interval estimate of θ . Important models include Gamma, Weibull, Fisher, among others. A familiar set-up where the results can be applied arises in random effects analysis of variance models with a Fisher distributed pivot (see Zhang and Woodroffe, 2002, for details). As in (A), for a sample $X = (X_1, \dots, X_n) \sim \frac{1}{\theta^n} f_1(\frac{x_1}{\theta}, \dots, \frac{x_n}{\theta}) \prod_i \mathbb{I}_{(0, \infty)}(x_i)$, Theorem 1 and Corollary 1 are applicable by conditioning on the maximal invariant statistic $V = (\frac{X_1}{X_n}, \dots, \frac{X_{n-1}}{X_n})$.

Remark 3. Further applications consist of power parameter families where we have a scale family for an observable Y and the model of interest are the distributions for $X = e^Y$. As a simple illustration, consider the Pareto model for X with densities $\frac{\gamma}{x^{\gamma+1}} \mathbb{1}_{(1, \infty)}(x)$ and the parametric constraint $\gamma \in (1, \gamma_0)$. In such cases, we have that $\gamma_0 \log(X) \sim \text{Exp}(\theta)$ with $\theta = \frac{\gamma_0}{\gamma} \geq 1$ and the results in (B) apply.

(C) **(location-scale families)** $(X_1, X_2) \sim \frac{1}{\theta_2^2} f_2(\frac{x_1 - \theta_1}{\theta_2}, \frac{x_2}{\theta_2}) \mathbb{I}_{(0, \infty)}(x_2)$; $\tau(\theta) = \theta_1 \geq 0$; $T(X, \theta) =$

$\frac{X_1 - \theta_1}{X_2}$; $\pi_H(\theta) = \frac{1}{\theta_2} \mathbb{I}_{(0, \infty)}(\theta_2) \mathbb{I}_{(-\infty, \infty)}(\theta_1)$, $\pi_0(\theta) = \frac{1}{\theta_2} \mathbb{I}_{(0, \infty)}(\theta_2) \mathbb{I}_{[0, \infty)}(\theta_1)$. This set-up encompasses, but is not limited to, the basic normal case: $Y_1, \dots, Y_n \sim^{ind.} N(\mu, \sigma^2)$ with σ^2 unknown and $\mu \geq 0$, and by taking X_1 and X_2 respectively as the sample mean and standard deviation of the Y_i 's. More generally, the results apply for linear models $Y = Z\beta + \epsilon$, $\epsilon \sim N(0, \sigma^2 I_n)$ where the objective is to estimate a lower-bounded linear combination $\tau(\theta) = l'\beta$, by setting $X_1 = \hat{\beta}(Z'Z)^{-1}Z'Y$, $X_2^2 = \|Y - Z\beta\|^2$, $\theta_1 = \beta$, $\theta_2 = \sigma$. Here, the pivot $T(X, \theta)$ has a Student distribution. Alternatively, if the objective is to estimate a lower bounded scale θ_2 , one can proceed as in (B).

(D) **(linear combination of several location parameters)** $X = (X_1, \dots, X_p) \sim f_0(x_1 - \theta_1, \dots, x_p - \theta_p)$; $\tau(\theta) = \sum_{i=1}^p a_i \theta_i$; $\pi_H(\theta) = \mathbb{I}_{\mathbb{R}^p}(\theta)$, $\pi_0(\theta) = \mathbb{I}_{[0, \infty)}(\tau(\theta))$, $T(X, \theta) = (\sum_{i=1}^p a_i X_i) - \tau(\theta)$. This set-up includes, for instance, estimating a difference $\theta_1 - \theta_2$ with an order constraint $\theta_1 \geq \theta_2$.

(E) **(multivariate location-scale families with homogeneous scale)**

In (D), we can incorporate a common scale and apply the results of this paper for estimating a lower bounded linear combination with $X = (X_1, \dots, X_p, X_{p+1}) \sim f_0(\frac{x_1 - \theta_1}{\theta_{p+1}}, \dots, \frac{x_p - \theta_p}{\theta_{p+1}}, \frac{x_{p+1}}{\theta_{p+1}})$, $\tau(\theta) = \sum_{i=1}^p a_i \theta_i$, $T(X, \theta) = \frac{(\sum_{i=1}^p a_i X_i) - \tau(\theta)}{X_{p+1}}$, and $\pi_0(\theta) = \frac{1}{\theta_{p+1}} 1_{(0, \infty)}(\theta_{p+1}) 1_{[0, \infty)}(\tau(\theta))$.

(F) **(several scale parameters)**

$(X_1, \dots, X_p) \sim (\prod_{i=1}^p \frac{1}{\theta_i}) f_1(\frac{x_1}{\theta_1}, \dots, \frac{x_p}{\theta_p})$; $\tau(\theta) = \sum_{i=1}^p a_i \log(\theta_i)$, $\pi_H(\theta) = \prod_i \frac{1}{\theta_i} \mathbb{I}_{(0, \infty)}(\theta_i)$, $\pi_0(\theta) = \pi_H(\theta) \mathbb{I}_{[0, \infty)}(\tau(\theta))$. This can consist, for instance with $p = 2$, $a_1 = 1$, $a_2 = -1$, of estimating a lower bounded ratio $\frac{\theta_2}{\theta_1} \geq 1$ of two scale parameters.

(G) **(quantiles in location-scale families)** $X_i \sim^{ind.} N(\mu, \sigma)$, $i = 1, \dots, n$, $\theta = (\mu, \sigma)$, $\tau(\theta) = \mu + \eta\sigma \geq 0$, $\pi_H(\theta) = \frac{1}{\theta_2} \mathbb{I}_{(0, \infty)}(\theta_2) \mathbb{I}_{(-\infty, \infty)}(\theta_1)$, $\pi_0(\mu, \sigma) = \frac{1}{\sigma} \mathbb{I}_{(0, \infty)}(\sigma) \mathbb{I}_{[0, \infty)}(\mu + \eta\sigma)$. $T(X, \theta) =$

$\frac{\bar{X} - \mu - \eta\sigma}{S}$. Here, $T(X, \theta)$ is distributed as non-central Student. The applications are not restricted to normality and are applicable in general for location-scale families as in (C).

6. Concluding remarks

For a large variety of situations with a lower bounded parametric constraint, we have obtained a class of Bayesian $(1 - \alpha) \times 100\%$ credible sets which provide minimal frequentist probability coverage exceeding $\frac{1-\alpha}{1+\alpha}$. These Bayesian confidence intervals include an equal tailed modification or approximation of the HPD credible set which coincides with the latter when the distribution of the underlying pivot is symmetric. In non-symmetric cases not covered by Marchand and Strawderman (2006), our findings provide instances of Bayesian credible sets with given minimal frequentist coverage and hence fill a gap in the literature. In comparison to earlier results for normal models, as well as the symmetric models considered by Marchand and Strawderman (2006), the findings here relative to the HPD are not new, but those related to other Bayesian credible sets are an addition. In seeking to evaluate the frequentist performance of Bayesian confidence intervals, our results illustrate that the choice of bounds or spending function matters, so that there does not necessarily exist a single universal assessment of their frequentist performance even in a given specific problem.

Acknowledgements

The authors are grateful to two reviewers, an associate editor, and the editor for useful comments and suggestions which led to a more self-contained and readable manuscript. Eric Marchand's research is supported in part by a grant from the Natural Sciences and Engineering Research Council of

Canada, and William Strawderman's research is partially supported by a grant from the Simons Foundation (#209035).

References

Berger, J.O. (1985). *Statistical Decision Theory and Bayesian Analysis*. Springer-Verlag, New York, 2nd edition.

Eaton, M. L. (1989). Group invariance Applications in Statistics. NSF-CBMS Regional Conference Series in Probability and Statistics 1. Hayward, CA.

Feldman, G.J. and Cousins, R. (1998). Unified approach to the classical statistical analysis of small signals. *Physical Review D*, **57**, 3873-3889.

Fraser, D.A.S. (2011). Is Bayes posterior just quick and dirty confidence? *Statistical Science*, **26**, 299-316.

Mandelkern, M. (2002). Setting Confidence Intervals for Bounded Parameters with discussion. *Statistical Science*, **17**, 149-172.

Marchand, É., Strawderman, W. E., Bosa, K., and Lmoudden, A. (2008). On the frequentist coverage of Bayesian credible intervals for lower bounded means. *Electronic Journal of Statistics*, **2**, 1028-1042.

Marchand, É. and Strawderman, W. E. (2006). On the behaviour of Bayesian credible intervals for some restricted parameter space problems. *Recent Developments in Nonparametric Inference and Probability : A Festschrift for Micheal Woodroffe*, IMS Lecture Notes-Monograph Series, **50**, pp. 112-126.

Robert, C.P. (2011). Discussion of "Is Bayes posterior just quick and dirty confidence?" by D.A.S. Fraser. *Statistical Science*, **26**, 317-318.

Roe, B. and Woodroffe, M. (2000). Setting confidence belts. *Physical Review D*, **63**, 013009/01-09.

Wasserman, L. (2011). Frasian inference. *Statistical Science*, **26**, 322-325.

Zhang, T. and Woodroffe, M. (2003). Credible and confidence sets for restricted parameter spaces. *Journal of Statistical Planning and Inference*, **115**, 479-490.

Zhang, T. and Woodroffe, M. (2002). Credible and confidence sets for the ratio of variance components in the balanced one-way model. *Sankhyā: Special issue in memory of D. Basu*, **64**, 545-560.